

AscoStabil

is a liquid feed supplement for poultry, rabbits and piglets to be used before, during and after stress exposure

Applications:

Stress exposure due to placement in stable, vaccinations and medication.

- Strengthens the immune system
- Reduces stress responses
- Lowers the morbidity and failure rate

Compositions:

Polyphenols, propylene glycol, sodium chloride

Ingredients:

Crude protein <0.3 %, crude fat 1.0 %, crude fibre 0.3 %, sodium 0.5 %, 3a300 Vitamin C 40g/l

Nutritional additives (per kg):

3a300 Vitamin C 40g/l

Feeding recommendation:

The use of MIRAVIT AscoStabil should be adapted to the vaccination schedule and the operating management in cooperation with the company veterinarian.

Broilers:

- Placement in stable: 5 days 1.5ml/l of drinking water
- Vaccination periods: 5 days 2ml/l of drinking water
- Stress (e.g. heat): 5 days 2ml/l of drinking water

Turkeys:

- Placement in stable: 5 days 1.5ml/l of drinking water
- from the 13th day after placement in stable: 5 days 2ml/l of drinking water

This product information is based on our current state of knowledge, experience and the statutory requirements. It describes the use of the product under standard operating conditions and assuming proper storage. We do not accept liability for any improper use and the resulting damage.

Free Hotline: Mr Berg 00490800 . 682-1133 or www.miravit.de

Breeding poultry:

- In the chicken coop: 5 days 2ml/l of drinking water during vaccination periods and stress

Parent animals: Chicken, ducks, geese, wild fowl:

- Female: 5 days 4ml/l of drinking water every 6th week
- Male: 1 day 2ml/l of drinking water every week
- Fattening parent animals: 5 days 2ml/l of drinking water every 3rd week up to the 28th week

Laying hens:

- 19 weeks old: 5 days 2ml/l of drinking water
- 25 weeks old: 5 days 2ml/l of drinking water

During intense stress (e.g. virus infections), take a 5-day break and then again 5 days 2ml/l of drinking water

Shelf life:

In the unopened, original packaging 12 months after production. Use within one month after opening the packaging

Store protected against light at 10° to 25°C. Reseal package after use.

Shake before use and offer diluted in water.

Packaging:

5 litre canister